

The Mosquitoes of Polynesia with a Pictorial Key  
to some Species Associated with Filariasis and/or  
Dengue Fever<sup>1,2</sup>

Yiau-Min Huang  
Medical Entomology Project  
Smithsonian Institution  
Washington, D.C. 20560

ABSTRACT. A list of the mosquitoes of Polynesia is tabulated and their distribution outlined. Keys for the identification of adults and larvae of Polynesian species are provided.

A pictorial key for the recognition of species associated with filariasis and dengue fever is furnished for the use of field workers.

#### INTRODUCTION

In order to assist field workers in recognizing the vector mosquitoes of filariasis and dengue in Polynesia, pictorial keys to the adult and larval stages have been prepared at the request of the World Health Organization. An attempt was made to make the keys precise, as simple as possible. A few additional characters indicated by a double asterisk (\*\*) have been added to certain species or species groups wherever necessary, to assure an exact identification and to avoid confusion with very similar and/or common species in the area. Unfortunately, the highly variable nature of the *scutellaris* group renders extremely difficult the identification of certain species of this group, some of which can only be identified by examination of the male terminalia. Therefore, it is always advisable that this examination be performed not only for routine confirmation of identification but also for the detection of new species in the area.

Map 1 shows the area of the South Pacific covered by the pictorial key. This area includes all of the Polynesian triangle north of the New Zealand faunal area. The 3 angles are represented by the Fiji Islands in the west, Easter Island in the east and the Hawaiian Islands in the north.

Table 1 lists all the 43 species and forms of mosquitoes known to occur

---

<sup>1</sup>Excluding the New Zealand faunal area covered by Belkin (1962).

<sup>2</sup>This study was supported by Research Contract No. DAMD-17-74-C-4086 from the United States Army Medical Research and Development Command, Office of the Surgeon General, Washington, D.C., and by a grant from the World Health Organization, Geneva, Switzerland.

# Report Documentation Page

Form Approved  
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>1977</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1977 to 00-00-1977</b>	
4. TITLE AND SUBTITLE <b>The Mosquitoes of Polynesia with a Pictorial Key to some Species Associated with Filariasis and/or Dengue Fever</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Medical Entomology Project, Smithsonian Institution, Washington, DC, 20560</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>see report</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

in the area delimited and shows by an asterisk (\*) the 19 known or suspected vector species included in the pictorial key. Those species having a limited distribution are also noted.

Table 2 lists the species by island or island group, indicating those that are endemic to a single area, and includes 9 new distribution records.

The non-pictorial keys to genera, subgenera, and species will obviate misidentification and will encourage interested workers to look for species not known to occur in these island groups and seek advice on the possible discovery of new species.

The keys will also assist in confirming the natural vectors of filariasis and/or dengue in the various island groups and in possibly incriminating species that are not known to be vectors at the present time.

It is important that final confirmation and/or determination of a species be made by specialists at one of the major museums such as the United States National Museum, Smithsonian Institution, Washington, D.C. 20560, or the Bernice P. Bishop Museum, P.O. Box 6037, 1355 Kalihi Street, Honolulu, Hawaii 96818. This is especially necessary for new distribution records, new vector species and possible new species.

Table 1  
Mosquito Species of Polynesia<sup>1</sup>

1. *Uranotaenia colocasiae* Edwards (Fiji Is. only)
2. *Uranotaenia painei* Edwards (Fiji Is. only)
- \*3. *Culex (Culex) quinquefasciatus* Say
4. *Culex (Culex) atriceps* Edwards (Society Is. only)
5. *Culex (Culex) kesseli* Belkin (Society Is. only; rare)
6. *Culex (Culex) marquesensis* Stone and Rosen (Marquesas Is. only)
7. *Culex (Culex) roseni* Belkin (Society Is. only)
8. *Culex (Culex) sitiens* Wiedemann
- \*9. *Culex (Culex) annulirostris* Skuse
10. *Culex (Culex) albinervis* Edwards (Fiji Is. only)
11. *Culex (Culex) samoensis* (Theobald) (Western Samoa only; rare)
12. *Aedeomyia (Aedeomyia) catasticta* Knab (Fiji Is. only)
13. *Mansonia (Coquillettidia) fijiensis* Belkin (Fiji Is. and rare in Western Samoa)
14. *Aedes (Finlaya) burnetti* Belkin (Fiji Is. only)
- \*15. *Aedes (Finlaya) fijiensis* Marks (Fiji Is. only)
16. *Aedes (Finlaya) freycinetiae* Laird (Fiji Is. only)
- \*17. *Aedes (Finlaya) oceanicus* Belkin
- \*18. *Aedes (Finlaya) samoanus* (Gruenberg) (Samoa Is. only)
- \*19. *Aedes (Finlaya) tutuilae* Ramalingam and Belkin (see Ramalingam and Belkin, 1965) (Samoa Is. only)
20. *Aedes (Finlaya) sp.* Albino form (Fiji Is. only)

21. *Aedes (Levua) suvae* Stone and Bohart (Fiji Is. only)
22. *Aedes (Ochlerotatus) edgari* Stone and Rosen (Society Is. only)
- \*23. *Aedes (Ochlerotatus) vigilax* (Skuse) (Fiji Is. only)
24. *Aedes (Aedimorphus) vexans* (Meigen)
- \*25. *Aedes (Stegomyia) aegypti* (Linnaeus)
- \*26. *Aedes (Stegomyia) albopictus* (Skuse) (Hawaiian Is. only)
- \*27. *Aedes (Stegomyia) cooki* Belkin (Niue I. and Tonga Is. only)
- \*28. *Aedes (Stegomyia) futunae* Belkin (Horn Is. only)
- \*29. *Aedes (Stegomyia) horrescens* Edwards (Fiji Is. only)
- \*30. *Aedes (Stegomyia) polynesiensis* Marks (see Huang, 1975)
- \*31. *Aedes (Stegomyia) pseudoscutellaris* (Theobald) (see Huang, 1975) (Fiji Is. only)
- \*32. *Aedes (Stegomyia) rotumae* Belkin (Rotuma I. only)
- \*33. *Aedes (Stegomyia) tabu* Ramalingam and Belkin (see Ramalingam and Belkin, 1965) (Tonga Is. only)
- \*34. *Aedes (Stegomyia) tongae* Edwards (Tonga Is. only)
- \*35. *Aedes (Stegomyia) upolensis* Marks (Samoa Is. only)
- \*36. *Aedes (Stegomyia) sp.* Tafahi form (Tonga Is. only)
37. *Aedes (Stegomyia) sp.* Wallis form (Wallis Is. only; rare)
38. *Tripteroides (Tripteroides) purpuratus* (Edwards) (Fiji Is. only)
39. *Tripteroides (Rachionotomyia) rotumanus* (Edwards) (Rotuma I. only)
40. *Toxorhynchites (Toxorhynchites) amboinensis* (Doleschall) (see Steffan, 1968) (introduced)
41. *Toxorhynchites (Toxorhynchites) brevipalpis* Theobald (see Steffan, 1968) (introduced)
42. *Toxorhynchites (Toxorhynchites) inornatus* (Walker) (introduced)
43. *Toxorhynchites (Toxorhynchites) splendens* (Wiedemann) (introduced)

---

<sup>1</sup>Excluding the New Zealand faunal area covered by Belkin (1962).

\*Included in the Pictorial Key.

Table 2  
Distribution of Mosquito Species in Polynesia<sup>1,2</sup>

1. Fiji Islands 23 spp.  
1, 2, 3, 8, 9, 10, 12, 13, 14, 15, 16, 20, 21, 23, 24, 25, 29, 30, 31, 34,\* 38, 42, 43
2. Tonga Islands 10 spp.  
3, 8, 9, 17, 24, 25, 27,\* 33, 34, 36
3. Samoa Islands 13 spp.  
3, 8, 9, 11 (rare), 13 (rare), 17, 18, 19, 24, 25, 30, 35, 41
4. Rotuma Island 6 spp.  
3,\*\* 9, 24, 32, 39, 43\*\* (no *Aedes aegypti*; *Culex sitiens* may be present)
5. Horn Islands 6 spp.  
3,\* 8,\* 9,\* 17,\* 28, 30 (no *Aedes aegypti*)
6. Wallis Islands 8 spp.  
3, 8, 9, 17,\* 24, 25, 30, 37
7. Ellis Islands 6 spp.  
3, 8, 9, 24, 25, 30
8. Tokelau Islands 2 spp.  
24, 30
9. Phoenix Islands (unknown)  
3??, 25??, 30?
10. Niue Island 4 spp.  
3, 8, 25, 27
11. Northern Cook Islands 4 spp.  
3, 9, 25, 27
12. Southern Cook Islands 5 spp.  
3, 9, 24, 25, 30
13. Society Islands 8 spp.  
3, 4, 5, 7, 9, 22, 25, 30
14. Austral Islands 4 spp.  
3, 9, 25, 30

- 15. Tuamotu Archipelago 4 spp.  
3, 9, 25, 30 (*Aedes albopictus* was introduced on one small island, but apparently did not become established)
- 16. Rapa Island 2 spp.  
3, 9
- 17. Pitcairn Island 3 spp.  
3, 9, 30
- 18. Easter Island 1 sp.  
3
- 19. Marquesas Islands 3 spp.  
3, 6, 30
- 20. Line Islands (unknown)  
3?, 25?, 30?
- 21. Hawaiian Islands 6 spp.  
3, 24, 25, 26, 40, 41 (There were no mosquitoes in the Hawaiian Islands until 1898 when a Spanish ship introduced the first species.)

<sup>1</sup>Excluding the New Zealand faunal area covered by Belkin (1962).

<sup>2</sup>Numbers under each island or island group correspond to the numbering of the mosquito species listed in Table 1; italicized numbers (e.g. 1, 15) indicate that the species is restricted to that island or island group.

\* New distribution records (J. C. Hitchcock, personal communication).

\*\* New distribution records (I. M. Rakai, personal communication to J.C. Hitchcock in "Report on mosquito survey - Rotuma", 17-19 August 1972).

KEYS TO GENERA, SUBGENERA, AND SPECIES IN POLYNESIA<sup>1</sup>

A. Adults

- 1. Apical half of proboscis bent sharply downward and backward and conspicuously more slender than basal half; posterior margin of scutellum evenly rounded . . . . . *Toxorhynchites*
- Apical half of proboscis not sharply bent downward and backward; posterior margin of scutellum distinctly trilobed . . . . . 2

- 2(1). Cell  $R_2$  always shorter than vein  $R_{2+3}$ ;  
wing membrane without distinct microtrichia . . . . . *Uranotaenia*
- Cell  $R_2$  always at least as long as vein  $R_{2+3}$ ;  
wing membrane with distinct microtrichia . . . . . 3
- 3(2). Spiracular setae present . . . . . 4
- Spiracular setae absent . . . . . 5
- 4(3). Vertex of head with azure  
blue scales . . . . . *Tripteroides (Tripteroides)*  
*purpuratus (Edwards)*
- Vertex of head without  
azure blue scales . . . . . *Tripteroides (Rachionotomyia)*  
*rotumanus (Edwards)*
- 5(3). Flagellomeres 12 and 13 relatively  
short and thick . . . . . *Aedeomyia (Aedeomyia)*  
*catastieta Knab*
- Flagellomeres 12 and 13 normal,  
neither short nor thick . . . . . 6
- 6(5). Postspiracular setae usually absent . . . . . 7
- Postspiracular setae usually present . . . . . 8
- 7(6). Claws of hindleg very small and inconspicuous;  
pulvilli present on all legs . . . . . *Culex*
- Claws of hindleg quite large and  
conspicuous; pulvilli absent . . . . . *Mansonia (Coquillettidia)*  
*fijiensis Belkin*
- 8(6). Wing scales broad; wings spotted; scutellum  
with broad scales on all lobes . . . . . *Aedes (Finlaya) kochi* group
- Wing scales narrow . . . . . 9
- 9(8). Head with decumbent scales largely broad, erect  
forked scales not numerous, restricted to occiput. *Aedes (Stegomyia)*
- Head with decumbent scales largely narrow, erect  
forked scales numerous, not restricted to occiput . . . . . 10
- 10(9). Pleural scaling restricted to posterior  
pronotum and sternopleuron . . . . . *Aedes (Levua) swae*  
Stone and Bohart

- Pleural scaling not restricted to posterior pronotum and sternopleuron . . . . . 11
- 11(10). Lower prealar scale patch present . . . . . *Aedes (Ochlerotatus)*
- Lower prealar scale patch absent . . . . . *Aedes (Aedimorphus) vexans* (Meigen)

---

<sup>1</sup>Excluding the New Zealand faunal area covered by Belkin (1962).

*Aedes (Finlaya) kochi* group

- 1. Tibiae with contrasting dark and light scales . . . . . 2
- Tibiae with all whitish scales . . . . . Fiji albino form
- 2(1). Halter largely dark scaled . . . . . *fijiensis* Marks
- Halter largely pale, yellow scaled . . . . . 3
- 3(2). Hind tarsomere 4 with all dark scales . . . . . *burnetti* Belkin
- Hind tarsomere 4 with at least some yellow scales ventrally or white scales apically . . . *freycinetiae* Laird

*Aedes (Ochlerotatus)*

- Anterior pronotum, propleuron and paratergite with scales . . . . . *vigilax* (Skuse)
- Anterior pronotum, propleuron and paratergite without scales . . . . . *edgari* Stone and Rosen

*Culex (Culex)*

- 1. Lower mesepimeral setae present; proboscis without a distinct complete median light ring; tarsi without distinct light rings . . . . . 2
- Lower mesepimeral setae absent; proboscis with a distinct complete median light ring; tarsi with distinct basal or basal and apical light rings . . . . . 5



- 2(1). Abdominal tergites with transverse basal pale bands connecting basolateral pale spots on some segments . . . . . 3
- Abdominal tergites with basolateral pale spots not connected by transverse basal pale bands on any segment . . . . . 4
- 3(2). Female: ventral surface of proboscis extensively pale scaled; male: palpus with white scales on ventral surface of segments 4 and 5 . . . . . *quinquefasciatus* Say
- Female: ventral surface of proboscis uniformly dark; male: palpus without white scales on ventral surface of segments 4 and 5 . . . *marquesensis* Stone and Rosen
- 4(2). Dorsal surface of hindfemur with basal 0.4 or more white . . . . . *kesseli* Belkin
- Dorsal surface of hindfemur with basal 0.1 or less white . . . . . *atriceps* Edwards
- 5(1). Abdominal tergites without any indication of transverse pale bands . . . . . *samoensis* (Theobald)
- Abdominal tergites with complete transverse pale bands on some segments . . . . . 6
- 6(5). A more or less conspicuous patch of broad erect scales in front of supraalar bristles . . . . . *albinervis* Edwards
- No broad erect scales in front of supraalar bristles. . . . . 7
- 7(6). Foretibia usually with a line of small pale spots on anterior surface along dorsal row of bristles . . *annulirostris* Skuse
- Foretibia usually without any pale spots on anterior surface along dorsal row of bristles . . . . . 8
- 8(7). Midfemur usually with some pale speckling on anterior surface . . . . . *sitiens* Wiedemann
- Midfemur usually without pale speckling on anterior surface . . . . . *roseni* Belkin

*Uranotaenia*

- Propleuron with scales; vein  $R_2$  at most 0.6 length of vein  $M_{1+2}$  . . . . . *colocasiae* Edwards

Propleuron without scales; vein  $R_2$  at  
 least 0.80 length of vein  $M_{1+2}$  . . . . . *painei* Edwards

*Toxorhynchites* (*Toxorhynchites*)

Females

1. Fore tarsomere 1 largely dark scaled;  
 upper posterior pronotum largely with  
 bluish and purplish scales . . . . . *brevipalpis* Theobald
- Fore tarsomere 1 largely light scaled . . . . . 2
- 2(1). Upper posterior pronotum largely with  
 bluish green scales . . . . . *inornatus* (Walker)
- Upper posterior pronotum largely with white scales . . . . . 3
- 3(2). Lateral scale tuft of abdominal segment  
 VI largely yellow . . . . . *splendens* (Wiedemann)
- Lateral scale tuft of abdominal segment  
 VI largely pale . . . . . *amboinensis* (Doleschall)

Males

1. Lateral scale tuft of abdominal segment  
 VIII black; upper posterior pronotum  
 largely with white scales . . . . . *amboinensis* (Doleschall)
- Lateral scale tuft of abdominal segment  
 VIII orange yellow . . . . . 2
- 2(1). Flagellomere 1 with dark scales on mesal  
 surface; upper posterior pronotum largely  
 with bluish and purplish scales . . . . . *brevipalpis* Theobald
- Flagellomere 1 with light scales on mesal surface . . . . . 3
- 3(2). Abdominal tergum I with dark scales medially  
 and yellowish scales laterally . . . . . *splendens* (Wiedemann)
- Abdominal tergum I with greenish scales  
 medially and white scales laterally . . . . . *inornatus* (Walker)

## B. Larvae

1. Median dorsal valve of siphon long, fixed, and with serrated dorsal margin . . . . . *Mansonia (Coquillettidia) fijiensis* Belkin
- Median dorsal valve of siphon short, movable, and without serrated dorsal margin . . . . . 2
- 2(1). Siphon with more than one pair of subventral (1-S) tufts . . . . . 3
- Siphon with a single pair of subventral (1-S) tufts . . . . . 5
- 3(2). Siphon with acus . . . . . *Culex*
- Siphon without acus . . . . . 4
- 4(3). Comb scales arising from a sclerotized plate . . . . . *Tripteroides (Tripteroides) purpuratus* (Edwards)
- Comb scales free, not arising from a sclerotized plate . . . . . *Tripteroides (Rachionotomyia) rotumanus* (Edwards)
- 5(2). Abdominal setae in groups of 3-5 on large common sclerotized plates . . . . . *Toxorhynchites*
- Abdominal setae arising separately and without strong sclerotized plates . . . . . 6
- 6(5). Antenna greatly swollen from base to setae 2-4 A . . . . . *Aedeomyia (Aedeomyia) catasticta* Knab
- Antenna at most slightly swollen proximad of seta 1-A . . . . . 7
- 7(6). Maxillary suture of head capsule at most barely indicated on anterior margin, never reaching posterior tentorial pit . . . . . *Uranotaenia*
- Maxillary suture of head capsule always complete and reaching posterior tentorial pit . . . . . 8

- 8(7). Abdominal segment I with seta 12 present . . . . . 9  
 Abdominal segment I with seta 12 absent . . . . . 11
- 9(8). Ventral brush with 5 pairs of setae,  
 each seta with long basal stalk, all  
 arising from basal boss, without  
 distinct bars, and no precratal  
 tufts . . . . . *Aedes (Finlaya) kochi* group
- Ventral brush with 5-7 pairs of setae  
 on grid, and with 2-4 precratal tufts . . . . . 10
- 10(9). Saddle large, extending on lateral surface;  
 seta 1-X on or adjacent to saddle . . . . . *Aedes (Ochlerotatus)*
- Saddle small, restricted to dorsal surface;  
 seta 1-X distinctly removed from saddle . . . . . *Aedes (Levua) suvae*  
 Stone and Bohart
- 11(8). Ventral brush with 6-7 pairs of setae  
 on grid, and with 2-4 precratal tufts . . . . . *Aedes (Aedimorphus)*  
*vexans* (Meigen)
- Ventral brush with 4 or 5 pairs of  
 setae, and no precratal tufts . . . . . *Aedes (Stegomyia)*
- Aedes (Finlaya) kochi* group
1. Comb scales in middle of posterior  
 row without fringe . . . . . *burnetti* Belkin
- Comb scales in middle of posterior  
 row with fringe . . . . . 2
- 2(1). Comb scale with 1, 2 pairs of sharp denticles  
 on basal part, distal part flattened, slightly  
 expanded, rounded apically and fringed . . . . . *fijiensis* Marks
- Comb scale without basal denticles, with a  
 slender long stem and a broad spatulate  
 apex, and fringed . . . . . *freycinetiae* Laird

---

The larva of Albino form is unknown.

*Aedes (Ochlerotatus)*

Siphon index more than 2.0; seta 5-C  
usually with 3-5 branches . . . . . *edgari* Stone and Rosen

Siphon index less than 2.0; seta 5-C  
usually with 1-2 branches . . . . . *vigilax* (Skuse)

*Culex (Culex)*

1. Antenna less than 0.35 head length; setae  
4, 6-C placed far forward on head capsule . . . . . 2
- Antenna more than 0.40 head length; setae  
4, 6-C placed farther back on head capsule . . . . . 4
- 2(1). Seta 1-C thick, spiniform . . . . . *marquesensis* Stone and Rosen
- Seta 1-C very thin . . . . . 3
- 3(2). Pecten tooth usually with 1-2 strong  
basal denticles . . . . . *atriceps* Edwards
- Pecten tooth usually simple . . . . . *kesseli* Belkin
- 4(1). Seta 1-C markedly flattened, its apex  
rounded or irregular . . . . . 5
- Seta 1-C very slender or moderately thickened,  
its apex acuminate or filamentous . . . . . 6
- 5(4). Saddle complete . . . . . *sitiens* Wiedemann
- Saddle incomplete . . . . . *roseni* Belkin
- 6(4). Seta 1-III-VI poorly developed, usually  
shorter than seta 3-III-VI . . . . . *albinervis* Edwards
- Seta 1-III-VI well developed, usually  
longer than seta 3-III-VI . . . . . 7
- 7(6). Seta 1-C very slender, filamentous distally,  
usually very lightly pigmented . . . . . *quinquefasciatus* Say
- Seta 1-C thickened, not filamentous distally,  
usually very strongly pigmented . . . . . *annulirostris* Skuse

---

The larva of *samoensis* (Theobald) is unknown.

*Uranotaenia*

- Seta 9-M, T single . . . . . *colocasiae* Edwards  
 Seta 9-M, T multiple . . . . . *painei* Edwards


*Toxorhynchites (Toxorhynchites)*

1. Seta 2-II-VI usually attached  
 to large dorsal plate . . . . . *inornatus* Walker  
 Seta 2-II-VI free from large dorsal plate . . . . . 2
- 2(1). Seta 11-IV, V usually with 3-4 branches . . . *splendens* (Wiedemann)  
 Seta 11-IV, V usually single or double . . . . . 3
- 3(2). Seta 9-C with 2-4 branches; 12-C  
 with 3-5 branches . . . . . *brevipalpis* Theobald  
 Seta 9-C with 5 or more branches;  
 12-C with 6 or more branches . . . . . *amboinensis* (Doleschall)

MORPHOLOGICAL FEATURES USED IN IDENTIFICATION


A. ADULT

Fig.1



THORAX - LATERAL

Fig. 2









Fig. 3

### WING-DORSAL



### CLAWS

### HINDLEG





Fig. 4

LARVA


1-X abdominal segments  
0-15 setae number

TERMINAL SEGMENTS  
Lateral

Fig. 5




ANAL SEGMENT (X)



*Mansonia (Coquillettidia)*

SIPHON





*Culex*

SIPHON


PICTORIAL KEY

A. ADULTS


Postspiracular setae absent.                      Postspiracular setae present.



(1) Lower mesepimeral setae present.



(1) Lower mesepimeral setae absent.



Culex quinquefasciatus Say

\*\*  
(2) Foretibia usually with a line of small pale spots on anterior surface along dorsal row of bristles.




Culex annulirostris Skuse

Erect forked scales numerous, not restricted to occiput.




Page 308

Erect forked scales not numerous, restricted to occiput.




Page 310

- (1) Scutellum with all narrow scales.
- (2) ♀ cercus long and slender.




- \*\*
- (3) Lower prealar scale patch present.
- (4) Subspiracular area without scales.




*Aedes vigilax* (Skuse)

- (1) Scutellum with all broad scales.
- (2) ♀ cercus short and broad.



- \*\*
- (3) Dorsal wing scales in contrasting pattern of dark and light scales.





- Pale areas
- a - Basal
  - b - Prehumeral
  - c - Humeral
  - d - Sectoral
  - e - Subcostal
  - f - Accessory subcostal

- Dark spots
- 1. basal
  - 2. prehumeral
  - 3. subbasal
  - 4. median
  - 5. preapical
  - 6. apical

(1) Accessory subcostal pale area not developed on vein C and  
(2) poorly developed, small on vein R1.

(1) Accessory subcostal pale area well developed, extensive on vein C as well as  
(2) on vein R1.




Prehumeral pale area of vein C developed.




Aedes fijiensis Marks

Prehumeral pale area of vein C not developed.




Aedes oceanicus Belkin

Sectoral pale area of vein C usually interrupted by a dark spot.




Aedes tutuilae Ramalingam & Belkin

Sectoral pale area of vein C usually not interrupted by a dark spot.



Aedes samoanus (Grunberg)

Scutum with lyre-shaped white markings.




Aedes aegypti (Linnaeus)

Scutum with a long median longitudinal white stripe extending from anterior margin to about level of wing root.



- (1) Abdominal tergites with complete basal transverse white bands and
- (2) with separate basolateral white spots.




lateral aspect

dorsal aspect

Aedes albopictus (Skuse)

- (1) Abdominal tergites without transverse white bands or with complete or incomplete subbasal white bands
- (2) connected to lateral white spots.




lateral aspect

dorsal aspect


↓  
Page 311

- (1) Supraalar white line more or less complete, with only narrow scales over wing root;
- (2) midlobe of scutellum with all broad white scales and without dark scales apically.




*Aedes futunae* Belkin

- (1) Supraalar white line complete, with broad flat scales over wing root;
- (2) midlobe of scutellum with broad white scales and with dark scales apically.



Lower mesepimeral white scale patch absent or very small, with no more than 3 scales.




Hind tarsomere 4 with basal 0.75 or more white.



*Aedes rotumae* Belkin

Lower mesepimeral white scale patch well developed, with at least more than 3 scales.



Hind tarsomere 4 with basal 0.60-0.70 white.






*Aedes upolensis* Marks

→ Page 312



Lateral prescutal white line present, or at least with some narrow white scales on scutal angle area.


Lateral prescutal white line not present.




Aedes pseudoscutellaris (Theobald)

Subspiracular area with scales.

Subspiracular area without scales.



Aedes horrescens Edwards





- Aedes polynesiensis Marks
- Aedes tongae Edwards
- Aedes tabu Ramalingam & Belkin
- Aedes cooki Belkin
- Aedes sp. Tafahi form

B. LARVAE

Siphon with more than 3 pairs of subventral tufts.



Siphon with a single pair of subventral tufts.



Page 314



(1) Mental plate usually with 6-9 teeth on each side of median tooth.

(1) Mental plate with at least 10 teeth on each side of median tooth.





(2) Seta 1-C thickened, not filamentous distally, usually very strongly pigmented.

(2) Seta 1-C very slender, filamentous distally, usually very lightly pigmented.



(3) Siphon with 5-7 pairs of subventral tufts.


(3) Siphon with 4 pairs of subventral tufts.




Culex annulirostris Skuse

Culex quinquefasciatus Say


- (1) Ventral brush with 6-7 pairs of setae and
- (2) with 2-4 precratal tufts.




- (1) Ventral brush with 4 or 5 pairs of setae and
- (2) no precratal tufts.




- \*\*
- (3) Comb in an irregular 2-3 rows,
  - (4) comb scale small, strongly fringed and usually with a differentiated apical spicule.
  - (5) Setae 1-VIII and 2-VIII not on common basal plate.



Aedes vigilax (Skuse)




- (1) Comb in a patch of several rows of scales, those of distal row elongate and varied in development.      (1) Comb in a single row.



- \*\*  
(2) Ventral brush with 5 pairs of setae, each seta with long basal stalk, all arising from basal boss, without distinct bars.



↓  
Page 316



↓  
Page 317

Seta 4-C usually with 2-5 branches, without enlarged base.

Seta 4-C usually at least with 6 branches, and always with enlarged base.







(1) Seta 4-C distinctly caudad of 6-C.

(1) Seta 4-C at about level, or slightly cephalad of 6-C.

(1) seta 4-C usually no more than 12 branches; 6-C usually with 3-7 branches.

(1) Seta 4-C at least 13 branches; 6-C usually with 8-12 branches.







(2) Siphon with uniform, dense, short spicules.

(2) Siphon largely bare or with short spicules in patches. (2)

(2) Siphon with uniform, short spicules. (2)

(2) Siphon with longer, denser and branched spicules. (2)




Aedes fijiensis Marks


Aedes oceanicus Belkin

Aedes tutuilae Ramalingam & Belkin

Aedes samoanus (Grunberg)




\*\*  
(2) Comb scale with very strong denticles at base of apical spine.




Aedes aegypti (Linnaeus)

Saddle complete.


Saddle incomplete.




Page 318



Page 319







Pecten tooth with very strong basal anterior denticles.



Aedes horrescens Edwards

Pecten tooth with rather small basal anterior denticles.




Aedes rotumae Belkin

Footnote : Adults

- Page 307 Culex sitiens Wiedemann, a widely distributed non-vector, is easily confused with Culex annulirostris, but does not have pale spots on the anterior surface of the foretibia (2).
- Page 308 Aedes vexans (Meigen) [=Aedes nocturnus (Theobald)], a widely distributed non-vector, is easily confused with Aedes vigilax, but does not have a scale patch at (3) and does not have a scale patch at (4). Aedes vigilax is only known from Fiji in Polynesia but is the major vector of subperiodic filariasis in New Caledonia and the Loyalty Islands.
- Page 311 Aedes sp. Tafahi form which could easily be confused with Aedes upolensis especially when the lower mesepimeral white scale patch is absent, has the dorsal surface of hind-femur with basal 0.25 or more white while in Aedes upolensis the dorsal surface of the hindfemur has basal 0.12 or less white. Aedes sp. Tafahi form is only known from Tonga.
- Larvae
- Page 313 Culex sitiens Wiedemann has seta 1-C thickened but it is irregularly dorsoventrally flattened and usually blunt while in Culex annulirostris it is thickly tapering and pointed. The 2 species are often found associated in the same breeding site.
- Page 314 Aedes vexans (Meigen) [=Aedes nocturnus (Theobald)] which could easily be confused with Aedes vigilax, has a single row of comb scales (3), a lightly fringed comb scale with a long apical spine (4), and setae 1-VIII and 2-VIII are on a common basal plate (5).
- Page 318 Aedes sp. Wallis form which could easily be confused with Aedes polynesiensis, has the saddle incomplete. Aedes sp. Wallis form is only known from the Wallis Islands.

MAP 1. AREA OF THE SOUTH PACIFIC COVERED BY THE PICTORIAL KEY



## ACKNOWLEDGMENTS

I am grateful to Dr. Ronald A. Ward for his helpful assistance and valuable comments in connection with this work and also for a critical review of the manuscript. I extend my thanks to Mr. Vichai Malikul for preparing the drawings. I also wish to express my gratitude to Dr. John N. Belkin, Department of Biology, University of California, Los Angeles, for the loan of the South Pacific specimens and for reviewing the keys and to Dr. Botha de Meillon, Philadelphia, Pennsylvania, who stimulated interest for the conduct of this study.

## REFERENCES

- Belkin, J. N. 1962. The mosquitoes of the South Pacific (Diptera: Culicidae). Berkeley and Los Angeles, University of California Press, 2 vols., 608 and 412 p.
- \_\_\_\_\_. 1964. The adults and pupa of *Culex (C.) kesseli* from Tahiti and remarks on the *atriceps* group (Diptera: Culicidae). Ann. Entomol. Soc. Am. 57:236-9.
- Huang, Yiau-Min. 1975. A redescription of *Aedes (Stegomyia) pseudoscutellaris* (Theobald) with a note on the taxonomic status of *Aedes (Stegomyia) polynesiensis* Marks (Diptera: Culicidae). Mosq. Syst. 7: 87-101.
- Ramalingam, S. and J. N. Belkin. 1965. Mosquito studies (Diptera, Culicidae). III. Two new species of *Aedes* from Tonga and Samoa. Contr. Am. Entomol. Inst. (Ann Arbor), 1(4): 1-10.
- Steffan, W. A. 1968. Hawaiian *Toxorhynchites* (Diptera: Culicidae). Proc. Hawaii. Entomol. Soc. 20: 141-55.